




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/62, 15/86, C07K 14/075, A61K 48/00</b>		<b>A1</b>	(11) International Publication Number: <b>WO 96/26281</b>
			(43) International Publication Date: 29 August 1996 (29.08.96)
(21) International Application Number: <b>PCT/US96/01957</b>		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AZ, BY, KG, KZ, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 13 February 1996 (13.02.96)		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: 08/395,381      21 February 1995 (21.02.95)      US			
(71) Applicants: GENVEC, INC. [US/US]; 12111 Parklawn Drive, Rockville, MD 20852 (US). CORNELL RESEARCH FOUNDATION, INC. [US/US]; Cornell Business and Technology Park, Suite 105, 20 Thornwood Drive, Ithaca, NY 14850 (US).			
(72) Inventors: WICKHAM, Thomas, J.; 5 Charen Court, Potomac, MD 20854 (US). FALCK-PEDERSEN, Erik; 8714 Buena Vista Drive, Dobbs Ferry, NY 10522 (US). ROELVINK, Petrus, W.; Apr. 203, 18313 Lost Knife Circle, Gaithersburg, MD 28799-2333 (US). BRUDER, Joseph, T.; 20423 Lindos Court, Gaithersburg, MD 20879 (US). GALL, Jason; 420 E. 70th Street #8P, New York, NY 10021 (US). KOVESDI, Imre; 206 Rollins Avenue, Rockville, MD 20052 (US).			
(74) Agents: KILYK, John, Jr. et al.; Leydig, Voit & Mayer, Ltd., Suite 4900, Two Prudential Plaza, 180 North Stetson, Chicago, IL 60601-6780 (US).			
(54) Title: CHIMERIC ADENOVIRAL FIBER PROTEIN AND METHODS OF USING SAME			

**Knob: (Ad5) amino acids 405-592****Shaft: (Ad5) amino acids 1-404****Penton Base**

## (57) Abstract

A recombinant adenovirus comprising a chimeric fiber protein and a therapeutic gene, a method of gene therapy involving the use of such an adenovirus, and an adenoviral transfer vector for the generation of such a recombinant adenovirus are provided.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

CHIMERIC ADENOVIRAL FIBER PROTEIN  
AND METHODS OF USING SAME

Technical Field Of The Invention

5       The present invention relates to a recombinant adenovirus comprising a chimeric adenoviral fiber protein and the use of a recombinant adenovirus comprising a chimeric adenoviral fiber protein in gene therapy.

10       Background Of The Invention

Adenoviruses belong to the family Adenoviridae, which is divided into two genera, namely Mastadenovirus and Aviadenovirus. Adenoviruses are nonenveloped, regular icosahedrons 65-80 nm in diameter (Horne et al.,  
15   J. Mol. Biol., 1, 84-86 (1959)). The capsid is composed of 252 capsomeres of which 240 are hexons and 12 are pentons (Ginsberg et al., Virology, 28, 782-783 (1966)). The hexons and pentons are derived from three different viral polypeptides (Maizel et al., Virology, 36, 115-125  
20   (1968); Weber et al., Virology, 76, 709-724 (1977)). The hexon comprises three identical polypeptides of 967 amino acids each, namely polypeptide II (Roberts et al., Science, 232, 1148-1151 (1986)). The penton contains a penton base, which is bound to the capsid, and a fiber,  
25   which is noncovalently bound to and projects from the penton base. The fiber protein comprises three identical polypeptides of 582 amino acids each, namely polypeptide IV. The adenovirus serotype 2 (Ad2) penton base protein is an 8 X 9 nm ring-shaped complex composed of five  
30   identical protein subunits of 571 amino acids each, namely polypeptide III (Boudin et al., Virology, 92, 125-138 (1979)). Proteins IX, VI, and IIIa are also present in the adenoviral coat and are thought to stabilize the viral capsid (Stewart et al., Cell, 67, 145-154 (1991);  
35   Stewart et al., EMBO J., 12(7), 2589-2599 (1993)).

Once an adenovirus attaches to a cell, it undergoes receptor-mediated internalization into clathrin-coated endocytic vesicles of the cell (Svensson et al., J.

Virol., 51, 687-694 (1984); Chardonnet et al., Virology, 40, 462-477 (1970)). Virions entering the cell undergo a stepwise disassembly in which many of the viral structural proteins are shed (Greber et al., Cell, 75, 477-486 (1993)). During the uncoating process, the viral particles cause disruption of the cell endosome by a pH-dependent mechanism (Fitzgerald et al., Cell, 32, 607-617 (1983)), which is still poorly understood. The viral particles are then transported to the nuclear pore complex of the cell (Dales et al., Virology, 56, 465-483 (1973)), where the viral genome enters the nucleus, thus initiating infection.

An adenovirus uses two separate cellular receptors, both of which must be present, to efficiently attach to and infect a cell (Wickham et al., Cell, 73, 309-319 (1993)). First, the Ad2 fiber protein attaches the virus to a cell by binding to an, as yet, unidentified receptor. Then, the penton base binds to  $\alpha_v$  integrins, which are a family of a heterodimeric cell-surface receptors that mediate cellular adhesion to the extracellular matrix molecules fibronectin, vitronectin, laminin, and collagen, as well as other molecules (Hynes, Cell, 69, 11-25 (1992)), and play important roles in cell signaling processes, including calcium mobilization, protein phosphorylation, and cytoskeletal interactions (Hynes, supra).

The fiber protein is a trimer (Devaux et al., J. Molec. Biol., 215, 567-588 (1990)) consisting of a tail, a shaft, and a knob. The fiber shaft region is composed of repeating 15 amino acid motifs, which are believed to form two alternating b-strands and b-bends (Green et al., EMBO J., 2, 1357-1365 (1983)). The overall length of the fiber shaft region and the number of 15 amino-acid repeats differ between adenoviral serotypes. For example, the Ad2 fiber shaft is 37 nm long and contains 22 repeats, whereas the Ad3 fiber is 11 nm long and contains 6 repeats. The receptor binding domain of the

fiber prot in is localized in the knob region encoded by the last 200 amino acids of the protein (Henry et al., J. of Virology, 68(8), 5239-5246 (1994)). The regions necessary for trimerization are also located in the knob region of the protein (Henry et al. (1994), supra). A deletion mutant lacking the last 40 amino acids does not trimerize and also does not bind to penton base (Novelli et al. Virology, 185, 365-376 (1991)). Thus, trimerization of the fiber protein is necessary for penton base binding. Nuclear localization signals that direct the protein to the nucleus to form viral particles following its synthesis in the cytoplasm are located in the N-terminal region of the protein (Novelli et al. (1991), supra). The fiber, together with the hexon, are the main antigenic determinants of the virus and also determine the serotype specificity of the virus (Watson et al., J. Gen. Virol., 69, 525-535 (1988)). The fiber protein is glycosylated with single N-acetyl-glucosamine residues; however, the functional significance of the glycosylation remains unclear (Caillet-Boudin et al., Eur. J. Biochem., 184, 205-211 (1989)).

Over ten fiber proteins from different adenoviral serotypes have been sequenced, only to reveal a larger sequence diversity than that observed among other adenoviral proteins. For example, the knob regions of the fiber proteins from the closely related Ad2 and Ad5 serotypes are only 63% similar at the amino acid level (Chroboczek et al., Virology, 186, 280-285 (1992)), whereas their penton base sequences are 99% identical. Ad2 and Ad5 fiber proteins, however, both likely bind to the same cellular receptor, since they cross-block each other's binding. In contrast, Ad2 and Ad3 fibers are only 20% identical (Signas et al., J. of Virology, 53, 672-678 (1985)) and presumably bind to different receptors, since each fails to cross-block the other's binding (Defer et al., J. of Virology, 64(8), 3661-3673 (1990)). Ad3 fiber utilizes sialic acid as its receptor,

whereas Ad2 fiber does not. Pretreatment of cells with neuraminidase or periodate abrogates Ad3, but not Ad2, binding. Also, soluble analogues of sialic acid block Ad3, but not Ad2, binding. However, sequence comparisons of the Ad2 and Ad3 fiber genes show distinct regions of conservation. Most of these regions are also conserved in the other human adenoviral fiber genes. Nonhuman adenoviral fiber genes show less homology to human serotypes but still trimerize. The receptors used by nonhuman serotypes are unknown. Recombinant adenoviral vectors have been used for the cell-targeted transfer of one or more recombinant genes to diseased cells or tissue in need of treatment. Such vectors are characterized by the advantage of not requiring host cell proliferation for expression of adenoviral proteins (Horwitz et al., *In Virology*, Raven Press, New York, vol. 2, pp. 1679-1721 (1990)); and Berkner, *BioTechniques*, 6, 616 (1988)), and, if the targeted tissue for somatic gene therapy is the lung, these vectors have the added advantage of being normally tropic for the respiratory epithelium (Straus, *In Adenoviruses*, Plenum Press, New York, pp. 451-496 (1984)).

Other advantages of adenoviruses as potential vectors for human gene therapy are as follows:

- (i) recombination is rare;
- (ii) there are no known associations of human malignancies with adenoviral infections despite common human infection with adenoviruses;
- (iii) the adenoviral genome (which is a linear, double-stranded DNA) can be manipulated to accommodate foreign genes that range in size;
- (iv) an adenoviral vector does not insert its DNA into the chromosome of a cell, so its effect is impermanent and unlikely to interfere with the cell's normal function;
- (v) the adenovirus can infect non-dividing or terminally differentiated cells, such as cells in the brain and lungs; and
- (vi) live adenovirus, having as an essential

characteristic the ability to replicate, has been safely used as a human vaccine (Horwitz et al. (1990), supra; Berkner et al. (1988), supra; Straus et al. (1984), supra; Chanock et al., JAMA, 195, 151 (1966); Haj-Ahmad et al., J. Virol., 57, 267 (1986); and Ballay et al., EMBO, 4, 3861 (1985)).

A drawback to adenovirus-mediated gene therapy is that significant decreases in gene expression are observed after two weeks following administration of the vector. In many therapeutic applications the loss of expression requires re-administration of the viral vector to overcome losses in expression. However, following administration of the viral vector, neutralizing antibodies are raised against both the fiber and hexon proteins (Wohlfart, J. Virology, 62, 2321-2328 (1988); Wohlfart et al., J. Virology, 56, 896-903 (1985)). This antibody response against the virus then can prevent effective re-administration of the viral vector. Accordingly, recombinant adenoviral vectors capable of avoiding such neutralizing antibodies that would allow repeated doses of adenoviral vectors to be administered in the context of gene therapy would represent a significant advance in current gene therapy methodology.

Another drawback of using recombinant adenovirus in gene therapy is that all cells that express the aforementioned two receptors used by adenovirus to attach and infect a cell will internalize the gene(s) being administered — not just the cells in need of therapeutic treatment. Likewise, certain cells, such as lymphocytes, which lack the  $\alpha_v$  integrin adenoviral receptors, are impaired in the uptake of adenoviruses (Silver et al., Virology 165, 377-387 (1988); Horvath et al., J. of Virology, 62(1), 341-345 (1988)) and are not readily amenable to adenovirus-mediated gene delivery. Accordingly, limiting adenoviral entry to specific cells or tissues and/or expanding the repertoire of cells amenable to adenovirus-mediated gene therapy would be a

significant improvement over the current technology.

Targeted adenoviral gene delivery should expand the cells amenable to gene therapy, reduce the amount of adenoviral vector that is necessary to obtain gene expression in the targeted cells, and reduce side effects and complications associated with increasing doses of adenovirus, such as inflammation and the transfection of normal, healthy cells.

Attempts have been made to target a virus to specific cells by sterically blocking adenoviral fiber protein with antibodies and chemically linking tissue-specific antibodies to the viral particle (Cotten et al., Proc. Natl. Acad. Sci. USA, 89, 6094-6098 (1992)). Although this approach has demonstrated the potential of targeted gene delivery, the complexity and reproducibility of this approach present major hurdles blocking its application in clinical trials. The difficulties thus far encountered in targeting the virus by these methods involve the method of synthesis required, which is to make major alterations in the viral particles following their purification. These alterations involve additional steps that covalently link large molecules, such as polylysine, receptor ligands and antibodies, to the virus (Cotten (1992), supra; Wagner et al., PNAS USA, 89, 6099-6103 (1992)). The targeted particle complexes are not homogeneous in structure and their efficiency is sensitive to the relative ratios of viral particles, linking molecules, and targeting molecules used.

The present invention seeks to overcome at least some of the aforesaid problems of recombinant adenoviral gene therapy. It is an object of the present invention to provide recombinant adenoviral vectors capable of avoiding neutralizing antibodies upon repeat administration, thereby enabling the maintenance of recombinant gene expression at a therapeutically effective level. It is another object of the present



invention to provide a cell-specific/tissue-specific recombinant adenovirus so as to target gene therapy to selected cells/tissues, thereby reducing the amount of recombinant adenoviral vector administered and any side-effects/complications. A further object of the present invention is to provide means for modifying the viral particle at the level of gene expression, thus allowing viral particles to be purified by conventional techniques. Another object of the present invention is to provide a method of gene therapy involving the use of such a homogeneous adenovirus, without the need for additional chemical modifications of viral particles, such as psoralen inactivation, or the addition of molecules to the virus which permit the covalent linkage of additional molecules to the virus. These and other objects and advantages of the present invention, as well as additional inventive features, will be apparent from the following detailed description.

#### Brief Summary Of The Invention

The present invention provides a recombinant adenovirus comprising a chimeric fiber protein, which differs from the native (wild-type) fiber protein by the introduction of a nonnative amino acid sequence. The nonnative amino acid sequence allows the adenovirus to be targeted towards a protein, such as a receptor or a bi- or multi-specific protein, which is specific for binding to the nonnative amino acid sequence and a target receptor, by facilitating direct binding between the nonnative amino acid sequence and the protein, i.e., receptor or bi/multi-specific protein. Alternatively, the nonnative amino acid sequence facilitates proteolytic removal of the chimeric fiber protein to allow targeting of the adenovirus by means of another adenoviral coat protein, such as the penton base. The present invention also provides an adenoviral transfer vector, among others, comprising a recombinant fiber gene sequence for

the generation of a chimeric fiber protein, and a method of using a protein-specific recombinant adenovirus, which is specific for a given receptor or bi-/multi-specific protein and which comprises a therapeutic gene, in gene therapy.

#### Brief Description Of The Figures

Figure 1 is a diagram of the penton complex.

Figure 2 is a partial restriction map of the vector pGBS.59-100.

Figure 3 is a partial restriction map of the vector p193 Ad5 Nde I/Sal I.

Figure 4 is a partial restriction map of the vector pAcSG2.

Figure 5 is a partial restriction map of the vector p193 Ad5 FC (F-).

Figure 6 is a partial restriction map of the vector p193 FC (F2).

Figure 7 is a partial restriction map of the vector pGBS.59-100 (F2).

Figure 8 is a partial restriction map of the vector pAcSG2 (F2).

Figure 9 is a partial restriction map of the vector p193 FC (F3).

Figure 10 is a partial restriction map of the vector pGBS.59-100 (F3).

Figure 11 is a partial restriction map of the vector pAcSG2 (F3).

Figure 12 is a partial restriction map of the vector p193 FC (HSF:RGD).

Figure 13 is a partial restriction map of the vector pGBS.59-100 (HSF:RGD).

Figure 14 is a partial restriction map of the vector pAcSG2 (HSF:RGD).

Figure 15 is a diagram of the construction of the vector pAd70-100dlE3.fiber7.

Detailed Description Of The Invention

The present invention provides, among other things, a recombinant adenovirus comprising a chimeric fiber protein. The chimeric fiber protein comprises a nonnative amino acid sequence, in addition to or in place of a native fiber amino acid sequence, which allows the adenovirus to bind to a protein, such as a receptor, which is other than a receptor bound by the native fiber, and which is referred to herein as a "target receptor," or a bi-/multi-specific protein, such as an antibody or fragment thereof, e.g., domain, with binding specificity for the nonnative amino acid sequence and for a target receptor. In the absence of native fiber amino acid sequences that enable trimerization of the native or chimeric fiber protein, the nonnative amino acid sequence comprises one or more sequences that enable trimerization of the chimeric fiber protein, which preferably are not immediately adjacent to the sequence that is specific for the aforesaid different protein, e.g., target receptor or bi- or multi-specific protein. Alternatively, the chimeric fiber protein comprises a nonnative amino acid sequence, in addition to or in place of a native fiber amino acid sequence, which is recognized by a protease and is cleaved by the protease, effectively removing the chimeric fiber protein and thereby allowing targeting of the adenovirus by means of another adenoviral coat protein, such as the penton base.

By "nonnative amino acid sequence" is meant any amino acid sequence that is not found in the native fiber of a given serotype of adenovirus and which is introduced into the fiber protein at the level of gene expression. "Nonnative amino acid sequence" includes an amino acid sequence from an adenoviral serotype other than the serotype of the adenovirus with the chimeric fiber protein. (For example, an Ad3 fiber amino acid sequence or the entire Ad3 fiber expressed in an Ad5 chimeric fiber protein or in place of an Ad5 fiber protein,

r spectively, is a "nonnative amino acid sequence.") It also includes a protease recognition sequence, i.e., a sequence that is recognized and cleaved by a protease.

By "protein-specific amino acid sequence" is meant  
5 any nonnative amino acid sequence encoding a protein, protein domain or peptide, whether specifically bound by another protein or fragment thereof, and is meant to include an amino acid sequence that confers upon a  
10 chimeric fiber the ability to directly bind to a target receptor or class of target receptors, preferably a cell-specific or tissue-specific receptor, and an amino acid sequence that confers upon a chimeric fiber the ability to directly bind to a bi- or multi-specific protein, such as an antibody or fragment thereof, e.g., domain, which  
15 binds to a target receptor(s).

By "receptor" is meant a protein, including membrane-bound and soluble proteins, with high specific affinity for biologically active substances, such as hormones, antibodies, and enzymes.

20 By "chimeric fiber protein" is meant a fiber protein comprising a nonnative amino acid sequence, which comprises either a protein binding sequence or a protease recognition sequence, in addition to or in place of a native fiber amino acid sequence, which comprises a  
25 protein binding sequence. "Chimeric fiber protein" is intended to include a fiber protein of a serotype which differs from that of the adenovirus on which it is expressed, i.e., where the entire native fiber sequence is replaced with an entire nonnative fiber sequence.

30 Incorporation of a protein-specific amino acid sequence into a chimeric fiber molecule allows targeting through two or more separate proteins which are chemically or otherwise linked to make a bi- or multi-specific protein. One component of the bi- or multi-specific protein binds to the fiber chimera. The second  
35 component or additional components of the bi- or multi-specific protein recognize(s) one or more additional

target receptors. For example, a bi- or multi-specific protein can include a bispecific multichain or single chain antibody (Cook et al., J. Immunol. Methods, 171, 227-237 (1994); Spooner et al., Human Pathol., 25, 606-614 (1994)) in which one domain specifically binds an epitope on chimeric fiber protein and the other domain specifically binds a target receptor. The bispecific antibodies bind to the chimeric fiber proteins in a recombinant adenovirus with the target receptor-specific domains of the bispecific antibodies available for binding to a target receptor.

Preferably, the entire native fiber protein or native receptor binding sequence of the fiber protein has been replaced at the DNA level with a nonnative protein-specific amino acid binding sequence. Alternatively, the native receptor binding sequence in the fiber gene has been rendered inactive at the DNA level by mutation of the sequence, such as by insertional mutagenesis, for example, or rendered conformationally inaccessible in the fiber protein, such as by insertion of a DNA sequence into or adjacent to the adenoviral fiber gene wherein "gene sequence" refers to the complete fiber gene sequence as well as any lesser gene sequence that is capable of being expressed as a functional fiber protein. For insertional mutagenesis, the DNA sequence encoding the native receptor binding sequence is preferably inserted near the gene sequence encoding the sequence encoding the native receptor binding sequence within the fiber gene sequence, so as to move the gene fiber protein, the native receptor binding sequence is conformationally inaccessible for binding to a receptor. In the latter case, the inserted nonnative gene sequence that causes the conformational inaccessibility of the native receptor binding sequence in the fiber protein is preferably one that is specific for a target receptor or bi- or multi-specific protein. Such a recombinant adenovirus can be used, for example, to study receptor

binding, adenoviral attachment, and adenoviral infection in vitro or in vivo.

In a preferred embodiment of the present invention, the above-described recombinant adenovirus additionally comprises a gene or genes capable of being expressed in a cell to which the virus has attached or by which the virus has been internalized and preferably is one having therapeutic utility, e.g., corrective DNA, i.e., DNA encoding a function that is either absent or impaired, or a discrete killing agent, such as DNA encoding a cytotoxin that, for example, is active only intracellularly, or DNA encoding ribozymes or antisense molecules. Accordingly, the use of the term "therapeutic gene" is intended to encompass these and any other embodiments of that which is more commonly referred to as gene therapy and is known to those of skill in the art. The recombinant adenovirus can be used for gene therapy or to study the effects of expression of the gene in a given cell or tissue in vitro or in vivo.

The recombinant adenovirus comprising a chimeric fiber protein and the recombinant adenovirus that additionally comprises a gene or genes capable of being expressed in a particular cell can be generated by use of a viral transfer vector, preferably an adenoviral transfer vector, in accordance with the present invention. The viral transfer vector, preferably an adenoviral transfer vector, comprises a chimeric adenoviral fiber gene sequence. The chimeric fiber gene sequence comprises a nonnative gene sequence in place of a native fiber gene sequence that encodes a receptor binding sequence, which has been deleted, or in addition to a native receptor binding sequence, which has been mutated or rendered conformationally inaccessible in the expressed chimeric fiber protein as described above. The nonnative sequence renders the adenovirus specific for binding to a protein, e.g., a receptor or bi- or multi-specific protein, as described above and, in the absence

of native trimerization sequences, contains a sequence(s) which allows the chimeric protein to trimeriz .

Alternatively, the nonnative sequence comprises an entire fiber sequence from an adenovirus of a different

- 5 serotype, which is then expressed in place of or in conjunction with native fiber on a given adenovirus. In other words, either all of the fibers on a given chimeric serotype or some of the fibers are of the native serotype, whereas others are of a nonnative serotype.

- 10 Another alternative is that the nonnative sequence comprises one or more of a protease recognition sequence, which is cleaved by a protease, thereby effecting removal of the chimeric fiber and targeting of the recombinant adenovirus by means of the penton base or other coat
- 15 protein (see Figure 1 for diagram of penton complex). Based upon the high degree of structural similarity between the fiber molecules of the more than 41 human serotypes of adenovirus, it is expected that any one of the serotypes of human or nonhuman adenovirus may be used
- 20 as the source of the fiber gene. It is preferred, however, that one of the serotypes for which the fiber gene has been sequenced is used.

- Restriction sites are introduced into the fiber gene sequence; preferably, such restriction sites are
- 25 introduced into or flanking a native receptor binding sequence of the fiber gene sequence by a suitable method, such as PCR mutagenesis. Preferably, these restriction sites are not already present in the fiber gene. Such sites facilitate the removal or inactivation, such as by
- 30 sequence alteration, of the DNA sequence encoding the native receptor binding sequence in a given adenoviral genome, or the rendering of the native receptor binding sequence conformationally inaccessible, thereby altering or eliminating the ability of the fiber to bind to a
- 35 receptor normally bound by the fiber. A deleted native receptor binding sequence can be replaced with, or a mutated or conformationally inaccessible receptor binding

sequence can be accompanied by, a different DNA sequence, preferably a DNA sequence encoding specificity for binding to a protein, such as a receptor, preferably a cell-specific or tissue-specific receptor, or class of receptors, or to a bi- or multi-specific protein with specificity for a given receptor, for example. Unique restriction sites in the fiber gene of one adenoviral serotype can be used to replace regions of the native fiber gene with homologous regions of the fiber gene from another serotype. Such restriction sites can even be used to replace an entire native fiber sequence with a nonnative fiber sequence.

Preferably, the adenoviral vector is one into which any suitable nonnative amino acid sequence can be rapidly inserted. For example, unique Nde I and Bam HI restriction sites in p193 FC(F) can be used to insert receptor binding sequences from other fiber serotype genes. Alternatively, sequences also can be inserted into the fiber gene sequence without the need for unique restriction sites through PCR. Because a recombinant adenovirus can be created via ligation of recombinant sequences with viral DNA or via homologous recombination, the adenoviral vector preferably has either (1) unique restriction sites that allow ligation of a vector fragment with the complementing fragments of the remaining viral genomes, as described in Example 1, or (2) adequate lengths of DNA on either side of the protein-specific sequence that allow efficient homologous recombination with viral DNA, as described in Example 1. A preferred adenoviral vector is shown in Figure 10, which is a partial restriction map of such a vector. The adenoviral vector of Figure 10 was generated as described in Example 1.

DNA encoding short peptide sequences or protein domains capable of binding to a given protein, preferably a receptor or class of receptors, in particular cell- or tissue-specific receptor, is preferred for insertion into



the fiber gene sequence in which the native receptor binding sequence has been deleted, mutated, or rendered conformationally inaccessible. However, other DNA sequences, such as those that encode bi-/multi-specific protein recognition sequences, such as receptor-specific antibody domains and sequences that encode antigenic epitopes recognized by specific antibodies, also may be used to replace the native receptor binding sequence. The target receptor is optimally cell-specific or tissue-specific, and desirably is expressed only on those cells or tissues to be treated.

A non-native, unique protease site also can be inserted into the fiber gene sequence to target an adenovirus through the penton base or penton base chimeras. The protease site preferably does not affect fiber trimerization or receptor specificity of the fiber protein. The fiber chimera-containing particles are produced in standard cell lines, e.g., those currently used for adenoviral vectors. Following production and purification, the particles are rendered fiberless through digestion of the particles with a sequence-specific protease, which cleaves the fiber proteins and releases them from the viral particles to generate fiberless particles. For example, thrombin recognizes and cleaves at the amino acid sequence Val Pro Arg Gly Ser (TRINS) (SEQ ID NO: 8) (Stenflo et al., J. Biol. Chem., 257, 12280-12290 (1982)). Fiberless particles have been shown to be stable and capable of binding and infecting cells (Falgout et al., J. of Virology, 62, 622-625 (1992)). These resultant particles then can be targeted to specific tissues via the penton base or other coat protein.

The size of the DNA used to replace the native receptor binding sequence may be constrained, for example, by impeded folding of the fiber or improper assembly of the penton base/fiber complex.

Alternatively, recombinant adenovirus comprising chimeric fiber protein may be produced by the removal of the native knob region, which comprises receptor-binding and trimerization domains, of the fiber protein and its replacement with a nonnative trimerization domain and a protein-specific binding domain (Peteranderl et al., Biochemistry, 31, 12272-12276 (1992)). A recombinant adenovirus comprising a chimeric fiber protein also may be produced by point mutation in the knob region and the isolation of clones that are capable of trimerization but incapable of binding to the native receptor. In either case, and also with respect to the removal and replacement of the native receptor-specific binding sequence as described above, new protein binding domains may be added onto the C-terminus of the fiber protein or into exposed loops of the fiber protein by inserting the nucleic acid sequence encoding the binding domain into the fiber gene sequence at the appropriate position.

Irrespective of which method is used to introduce a protein binding sequence into the fiber protein, the fiber protein must be able to trimerize. If the fiber protein cannot trimerize, it will be unable to bind to penton base protein. Accordingly, the native receptor binding sequence must be changed without affecting the ability of the molecule to trimerize.

A recombinant chimeric fiber gene sequence can be moved from an adenoviral transfer vector into baculovirus or a suitable prokaryotic or eukaryotic expression vector for expression and evaluation of receptor or protein specificity and avidity, trimerization potential, penton base binding, and other biochemical characteristics. Accordingly, the present invention also provides recombinant baculoviral and prokaryotic and eukaryotic expression vectors comprising a chimeric adenoviral fiber gene sequence. The chimeric fiber gene sequence includes a nonnative sequence in addition to or in place of a native fiber amino acid sequence, which is specific for

binding to a protein other than a protein bound by the native fiber. The native fiber amino acid sequence may be deleted, mutated, or rendered conformationally inaccessible as described above with respect to the recombinant adenovirus comprising a chimeric fiber protein. By moving the chimeric gene from an adenoviral vector to baculovirus or a prokaryotic or eukaryotic expression vector, high protein expression is achievable (approximately 5-50% of the total protein being the chimeric fiber). Accordingly, the present invention also provides a recombinant baculovirus comprising a chimeric fiber gene and a chimeric adenoviral fiber protein comprising a nonnative amino acid sequence in addition to or in place of a native fiber amino acid sequence. The nonnative amino acid sequence is specific for binding to a protein, such as a receptor or a bi-/multi-specific protein, or encodes a protease cleavage site as described above. For protein characterization studies, the recombinant chimeric fiber protein (rcF protein, such as rcF5) can be purified using any suitable methods, such as those described by Wickham et al. (1993), supra.

Various characteristic parameters of the fiber protein of interest can be assessed. Specificity and affinity of the receptor or other protein/rcF interaction can be assessed by Scatchard analysis as shown previously by Wickham et al. (1993), supra, for wild-type penton base protein. Receptor specificity can be further assessed by using antibodies and peptides specific for the targeted receptor to block rcF5 binding to cells, using conventional methods. rcF binding to penton base protein can be assessed by its ability to precipitate radiolabeled penton base protein when coupled to protein A-coated beads via an antibody to the fiber protein.

Viral attachment, entry and gene expression are evaluated initially by using the adenoviral vector containing the insert of interest to generate a recombinant virus expressing the chimeric fiber protein

and a marker gene, such as  $\beta$ -galactosidase.  $\beta$ -galactosidase expression in cells infected with adenovirus containing the  $\beta$ -galactosidase gene (Ad-LacZ) can be detected as early as two hours after adding Ad-Gluc to cells. This procedure provides a quick and efficient analysis of cell entry of the recombinant virus and gene expression, and is implemented readily by an artisan of ordinary skill using conventional techniques.

A recombinant virus, which lacks a native receptor binding sequence in the fiber, can be produced in human embryonic cell line 293 (HEK 293), which allows replication of Ad5LacZ virus in which the LacZ gene replaces the E1 region of the adenoviral genome. For producing recombinant adenovirus containing chimeric fiber, the 293 cell line must express the receptor to which the chimeric fiber protein is targeted. In the absence of constitutive receptor expression, the receptor gene can be transfected into the 293 cell line to create a stably expressing cell line.

Recombinant adenoviruses of the present invention can be used to treat any one of a number of diseases by delivering to targeted cells corrective DNA, i.e., DNA encoding a function that is either absent or impaired, or a discrete killing agent, e.g., DNA encoding a cytotoxin that, for example, is active only intracellularly. Diseases that are candidates for such treatment include, for example, cancer, e.g., melanoma, glioma or lung cancers; genetic disorders, e.g., cystic fibrosis, hemophilia or muscular dystrophy; pathogenic infections, e.g., human immunodeficiency virus, tuberculosis or hepatitis; heart disease, e.g., preventing restenosis following angioplasty or promoting angiogenesis to reperfuse necrotic tissue; and autoimmune disorders, e.g., Crohn's disease, colitis or rheumatoid arthritis.

One skilled in the art will appreciate that suitable methods of administering a recombinant adenovirus of the present invention to an animal for purposes of gene

therapy (see, for example, Rosenfeld et al., Science,  
252, 431-434 (1991); Jaffe et al., Clin. Res., 39(2),  
302A (1991); Rosenfeld et al., Clin. Res., 39(2), 311A  
(1991); Berkner, BioTechniques, 6, 616-629 (1988)),  
5 chemotherapy, and vaccination are available, and,  
although more than one route can be used to administer  
such a recombinant adenovirus, a particular route can  
provide a more immediate and more effective reaction than  
another route. Pharmaceutically acceptable excipients  
10 are also well-known to those who are skilled in the art,  
and are readily available. The choice of excipient will  
be determined in part by the particular method used to  
administer the recombinant adenovirus. Accordingly,  
there is a wide variety of suitable formulations for use  
15 in the context of the present invention. The following  
methods and excipients are merely exemplary and are in no  
way limiting.

Formulations suitable for oral administration can  
consist of (a) liquid solutions, such as an effective  
20 amount of the compound dissolved in diluents, such as  
water, saline, or orange juice; (b) capsules, sachets or  
tablets, each containing a predetermined amount of the  
active ingredient, as solids or granules; (c) suspensions  
in an appropriate liquid; and (d) suitable emulsions.  
25 Tablet forms can include one or more of lactose,  
mannitol, corn starch, potato starch, microcrystalline  
cellulose, acacia, gelatin, colloidal silicon dioxide,  
croscarmellose sodium, talc, magnesium stearate, stearic  
acid, and other excipients, colorants, diluents,  
30 buffering agents, moistening agents, preservatives,  
flavoring agents, and pharmacologically compatible  
excipients. Lozenge forms can comprise the active  
ingredient in a flavor, usually sucrose and acacia or  
tragacanth, as well as pastilles comprising the active  
35 ingredient in an inert base, such as gelatin and  
glycerin, or sucrose and acacia, emulsions, gels, and the

like containing, in addition to the active ingredient, such excipients as are known in the art.

The recombinant adenovirus of the present invention, alone or in combination with other suitable components, can be made into aerosol formulations to be administered via inhalation. These aerosol formulations can be placed into pressurized acceptable propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like. They may also be formulated as pharmaceuticals for non-pressured preparations such as in a nebulizer or an atomizer.

Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain anti-oxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The formulations can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid excipient, for example, water, for injections, immediately prior to use. Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described.

Additionally, the recombinant adenovirus of the present invention may be made into suppositories by mixing with a variety of bases such as emulsifying bases or water-soluble bases.

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams, or spray formulas containing, in addition to the active ingredient, such carriers as are known in the art to be appropriate.

The dose administered to an animal, particularly a human, in the context of the present invention will vary with the genus of interest, the composition employed, the method of administration, and the particular site and organism being treated. However, the dose should be sufficient to effect a therapeutic response.

In addition to the recombinant adenovirus of the present invention, the recombinant vectors, e.g., the adenoviral transfer vector, also have utility in vitro. They can be used as a research tool in the study of adenoviral attachment and infection of cells and in a method of assaying receptor-ligand interaction. Similarly, the recombinant fiber protein comprising a nonnative amino acid sequence in addition to or in place of a native receptor binding sequence can be used in receptor-ligand assays and as adhesion proteins in vitro or in vivo, for example.

The following examples further illustrate the present invention and, of course, should not be construed as in any way limiting its scope.

#### Example 1

This example describes how to change adenoviral antigenicity without changing receptor specificity by creating a chimeric fiber protein in which the native Ad5 receptor binding domain is replaced with the nonnative Ad2 receptor binding domain.

The Ad2 fiber gene was amplified by PCR, wherein an Xho I site was incorporated into the 5' end of the sense PCR primer of SEQ ID NO:1, and Xma I and Bam HI sites were incorporated into the 5' end of the antisense primer of SEQ ID NO:2 to allow cloning into the Xho I/Xma I cloning sites in the vector pAcSG2 (Figure 4) (Pharmingen, San Diego, CA) to create the vector pAcSG2 (F2) (Figure 8). The pAcSG2 (F2) was used to evaluate

th fiber chimera at the protein level for receptor and penton base binding activity.

The Nde I/Bam HI fragment of the fiber2 gene was removed from pAcSG2 (F2) and cloned into the vector p193 FC (F-) (Figure 5) to create p193 FC (F2) (Figure 6).  
The vector p193 FC (F-) was used as the base vector for making all chimeric fiber adenoviruses. The p193 FC (F-) vector was created by cutting the p193 Ad5 (Nde I/Sal I) (Figure 3) vector with Nde I and Mun I to remove most of the Ad5 fiber gene, including its stop and polyadenylation signals, and by replacing the Nde I/Mun I fragment with a synthetic oligonucleotide, which lacks the amino acid coding region for Ad5 fiber but retains the Ad5 fiber stop and polyadenylation signal. The synthetic oligonucleotide was prepared from two sense and antisense complementary oligonucleotides, SEQ ID NO:3 and SEQ ID NO:4, respectively, which recreate cut Nde I and Mun I sites when paired and contain a Bam HI site just upstream of the stop codon to allow directional cloning into the Nde I/Bam HI sites. The Nde I/Sal I fragment containing the chimeric Ad2/Ad5 fiber gene was then cloned into the vector pGBS.59-100 (Figure 2) to create the transfer vector pGBS.59-100 (F2) (Figure 7). The pGBS.59-100 (F2) transfer vector was then cut with Sal I, purified and transfected into an appropriate cell line with a complementing 27,530 bp Ad5 DNA fragment (left arm, 0-27,530 bp) to create recombinant virus through homologous recombination. An appropriate cell line is any cell line which expresses the receptor for the chimeric fiber and which is capable of replicating the adenoviral vector. The complementing fragment of Ad5 DNA was prepared by cutting the Ad5 DNA with the restriction enzyme Srf I, which cuts the Ad5 genome once at position 27,530 in the wild-type Ad5 genome. The larger 27,530 bp piece was then isolated from the smaller bp fragment using a CsCl gradient, although an agarose



and AdCMV-CATNeo-dlE3-Fiber7 (AdCMV-CATNeo + pAd70-100dlE3.Fiber7) viruses. Each virus was grown in large scale, i.e., a 1 l suspension of human embryonic kidney 293 cells, to yield virus at a concentration of  $10^{12}$  particles/ml. A549 cells were infected with an estimated 100, 300 or 1,000 particles/cell of one of the three viruses. After 48 hr, the cells were harvested and lysates were prepared as described in Kass-Eisler et al., PNAS USA, 90, 11498-11502 (December 1993). Using 50  $\mu$ l of each lysate, CAT assays were performed and acetylated chloramphenicol products were separated by thin layer chromatography using chloroform:methanol (95:5). The results of the assays indicated that each virus was able to infect cells and express gene products at appropriate levels. Accordingly, the virus in which the native fiber was replaced with a nonnative fiber could infect cells and express genes like the parental virus.

Following this study, adult Sprague Dawley rats were infected with  $10^8$  viral particles by direct cardiac injection as described in Kass-Eisler et al., supra. Five days later, the rats were sacrificed, cardiac lysates were prepared, and CAT assays were performed. The amount of the CAT gene product produced was compared between the dlE3 and dlE3-Fiber 7 viruses. Results indicated that both viruses were able to infect cells in vivo. The replacement of the wild-type Ad5 fiber gene with that of Ad7 did not impair the ability of the virus to infect cells. Accordingly, the virus in which the native fiber was replaced with a nonnative fiber could also infect cells and express genes like the parental virus in vivo. These results support the utility of adenovirus with chimeric fiber in the context of gene therapy.

All publications cited herein are hereby incorporated by reference to the same extent as if each publication were individually and specifically indicated

gel or other appropriate separation technique could have been utilized.

Alternatively, viral DNA can be cut with a restriction enzyme, such as Spe I, which cuts at position 27,082 in the wild-type Ad5 genome. The 27,082 bp Spe I fragment can be isolated from the smaller fragment as described above and then ligated with the complementing Spe I/Sal I fragment from the pGBS.59-100 (F2) vector and then transfected into the appropriate cell line. Recombinant virus then can be isolated by plaque assay and verified as recombinant using PCR probes specific for the chimera and by restriction analysis.

#### Example 2

This example describes how to change receptor specificity and antigenicity by creating a chimeric fiber protein in which the native Ad5 receptor binding domain is replaced with the nonnative Ad2 receptor binding domain.

Oligonucleotide primers were used to amplify a large fraction of the Ad3 fiber gene using PCR. The 5' sense primer of SEQ ID NO:5 contained an in-frame mutation that incorporated an Nde I site, whereas the antisense oligonucleotide of SEQ ID NO:6 incorporated a Bam HI site to allow cloning of the amplified fragment into pAcSG2 (F2) in which the corresponding Nde I/Bam HI region of the Ad2 fiber gene was removed. The Nde I/Bam HI fragment of the gene for Ad3 fiber was then removed from the vector pAcSG2 (F3) (Figure 11) and cloned into the vector p193 FC (F-) to create p193 FC (F3) (Figure 9). The Nde I/Sal I fragment containing the chimeric Ad3/Ad5 fiber gene was then cloned into the vector pGBS.59-100 to create the transfer vector pGBS.59-100 (F3) (Figure 10). The pGBS.59-100 (F3) transfer vector was then cut, purified, and transfected into the appropriate cell line with Ad5 arms as in Example 1. Recombinant virus was

then isolated and verified to be recombinant as in  
Exempl 1.

The receptor for Ad3 contains a sialic acid  
component, which is required for binding of Ad3, while  
5 binding of Ad5 does not involve sialic acid. Since  
sialic acid is found on all higher eukaryotic cells, the  
Ad3/Ad5 fiber chimera is capable of binding to all cells.  
Such a vector can infect a broader range of cell types  
and exhibits different tissue specificity than non-  
10 chimeric Ad5 vectors in vivo.

### Example 3

This example describes how receptor specificity can  
be changed and binding domains can be incorporated at the  
15 C-terminus of mouse adenoviral fiber.

The fiber sequence from a nonhuman adenoviral  
serotype, mouse adenovirus type 1, for example, is  
amplified using PCR. Nhe I and Bam HI sites incorporated  
into the sense and antisense PCR primers, respectively,  
20 allow subsequent cloning of the PCR product. The Nhe I  
site corresponds to a naturally occurring site in Ad5  
fiber that occurs after the sequence encoding penton base  
recognition domains. The antisense primer, in addition  
to the required Bam HI site, contains a sequence encoding  
25 an  $\alpha_v\beta_3$ -specific RGD peptide following an amino acid  
spacer of 5-30 amino acids (such as poly [Ala Ser] or  
poly [Gly]). A unique restriction site is incorporated  
into the sequence following the spacer sequence and then  
again before the stop codon. The site allows the  
30 incorporation of receptor-specific sequences other than  
the  $\alpha_v\beta_3$ -specific RGD peptide. The resultant PCR product  
is then cloned into pAcSG2 (F5) to replace the  
corresponding Ad5 fiber sequence and create pAcSG2  
(MouseRGD). The Nde I/Bam HI fragment containing the  
35 chimeric fiber gene is cloned into p193 FC (F-) to create  
p193 FC (MouseRGD). The Nde I/Sal I fragment from p193  
FC (MouseRGD) is cloned into pGBS.59-100 to create the

transfer vector pGBS.59-100 (MouseRGD). The transfer vector is then prepared and transfected along with complementing Ad5 DNA into cells expressing the  $\alpha_v\beta_3$  receptor as described in Example 1. Recombinant virus  
5 containing the chimeric fiber gene is analyzed as in Example 1. Using the unique restriction site incorporated into the vector, other receptor binding domains, such as the P-selectin binding domain or a single chain receptor-specific antibody, can be directly  
10 cloned into the vector. However, the cell line used for transfection must express the targeted receptor in order for the recombinant virus to attach and infect cells. Incorporation of receptor or antibody binding domains into fiber molecules that do not recognize human  
15 receptors allow for the targeting of a vector using such a fiber without retaining residual amino acid sequences that recognize human receptors and prevent efficient targeting.

20

#### Example 4

This example describes how to change receptor specificity by mutating a native fiber receptor-binding domain and incorporating a nonnative binding domain at the C-terminus or within an exposed loop of a mutant Ad5.

25 A mutated fiber gene, one which generates fiber that can trimerize but cannot bind to a native fiber receptor, is amplified by PCR using primers that incorporate proper restriction sites for cloning. The antisense primer, in addition to the required Bam HI site, contains a sequence  
30 encoding an  $\alpha_v\beta_3$ -specific RGD peptide following an amino acid spacer of 5-30 amino acids, such as poly (Ala Ser) or poly Gly. A unique restriction site is incorporated into the sequence following the spacer sequence and then before the stop codon. The site allows the incorporation  
35 of receptor-specific sequences other than the  $\alpha_v\beta_3$ -specific RGD peptide. The amplified chimeric gene is cloned into the p193 FC (F-) plasmid to obtain p193 FC

(F5': $\beta$ 3). The Nde I/Sal I fragment containing the chimeric fiber gene is then cloned into the pGBS.59-100 vector to obtain pGBS.59-100 (F5': $\beta$ 3). The transfer vector is prepared and transfected with complementing Ad5 DNA as described in Example 1. Recombinant virus containing the chimeric fiber gene is analyzed as in Example 1. Other receptor-specific or antibody-specific binding domains can be cloned into the vector to create fiber chimeras with such sequences at the C-terminus of the protein or within exposed loops of the fiber molecule for targeting to other receptors or antibodies, respectively, as described in Example 3.

#### Example 5

15 This example describes replacement of a knob with a trimerization domain and the incorporation of a binding domain at the C-terminus of the knob protein.

The adenovirus type 2 fiber gene was amplified using PCR from Ad2 viral DNA and cloned into the baculovirus transfer vector, pBlueBac2 (Invitrogen, La Jolla, CA), to generate the vector pBB2F. The unique restriction sites Pst I and Bam HI encompass the region of the fiber2 gene encoding the knob region of the protein. These sites were used to remove the Pst I to Bam HI portion of the fiber gene and to replace it with DNA encoding the trimerization domain from the heat shock factor (HSF) protein of K. lactis fused via a glycine spacer to an RGD peptide specific for the integrin  $\alpha v \beta 3$ . The DNA encoding the HSF domain and RGD peptide was obtained through PCR from a plasmid containing the sequence for the entire K. lactis HSF protein. The DNA sequence encoding the RGD peptide was incorporated into the antisense DNA primer of SEQ ID NO:7 used in the PCR of the HSF trimerization domain to create the DNA sequence encoding the HSF:RGD fusion protein. The sense primer contained a Pst I site native to the Ad2 fiber gene. The PCR product was then digested with Pst I and Bam HI and cloned into the pAcSG2

(F2) vector to obtain the plasmid pAcSG2:HSF:RGD (Figure 14). Unique Spe I and Sca I sites were incorporated into the chimeric HSF:RGD gene so that different receptor-specific or antibody-specific sequences could be rapidly inserted into the gene in place of the RGD-coding sequence at the end of the sequence encoding the glycine spacer arm. The pAcSG2:HSF:RGD plasmid (Figure 14) was used to make recombinant baculovirus which expresses the fusion protein at high levels. The fusion protein expressed was the correct size and formed a trimer. The Nde I/Bam HI fragment of the chimeric gene was then removed from the vector pAcSG2 (HSF:RGD) and cloned into the vector p193 FC (F-) to create p193 FC (HSF:RGD) (Figure 12). The Nde I/Sal I fragment containing the chimeric fiber gene was cloned into the vector pGBS.59-100 (Figure 2) to create the transfer vector pGBS.59-100 (HSF:RGD) (Figure 13). The pGBS.59-100 (HSF:RGD) transfer vector was then cut, purified and transfected into the appropriate cell line with Ad5 arms as in Example 1. Recombinant virus was then isolated and verified to be recombinant as in Example 1.

#### Example 6

This example describes how to replace a knob with a trimerization domain and how to incorporate a binding domain containing a protease cleavage site at the C-terminus of the knob.

A chimeric fiber can be targeted to a new receptor by incorporating an epitope into the chimera which is recognized by a bi-specific antibody. An additional RGD domain is incorporated at the C-terminus of the protein and separated from the antibody epitope by a unique protease recognition site. The chimeric virus is capable of growing in tissue culture cells that express the receptor for the RGD sequence. Final preparations of virus are then exposed to the protease to remove the RGD sequence, leaving the epitope. The viral particles are

th n exposed to a bi-sp cific antibody in which on half  
of the molecule recognizes the epitope on the chimeric  
fiber and the other half recognizes any desired receptor,  
e.g., cell- or tissue-specific, receptor. The RGD  
5 sequence is absent and the virus binds to and enters only  
those cells recognized by the bi-specific antibody.

#### Example 7

This example describes how to change the adenoviral  
10 antigenicity and receptor specificity of an Ad5 virus by  
replacing native Ad5 fiber with nonnative Ad7 fiber and  
demonstrates the ability of such recombinant virus to  
infect cells in vitro and in vivo.

The Ad5 virus - Ad7 fiber construct was generated as  
15 shown in Figure 15. An approximately 2.7 kb (Ad5 28689-  
31317 bp) fragment in pAd70-100 was replaced with a Pac I  
linker (pAd70-100dlE3.Pac). A Bam HI linker was inserted  
at a unique Mun I site as indicated in Figure 13 to  
produce pAd70-100dlE3.Pac.Bam. A PCR-amplified Pac I-Bam  
20 HI fragment of approximately 1.1 kb containing the Ad7  
fiber gene was inserted into pAd70-100dlE3.Pac.Bam to  
produce pAd70-100dlE3.fiber7.

In order to assess the ability of Ad5 virus with Ad7  
fiber to infect cells in vitro and in vivo, reporter gene  
25 assays were performed. A replication-defective  
recombinant adenoviral reporter vector designated AdCMV-  
CATNeo was used in the reporter gene assay. The reporter  
vector consists of the adenoviral origin of replication  
and viral packaging sequences, a combination of strong  
30 eukaryotic promoter (cytomegalovirus or CMV-1) and  
splicing elements, the bacterial chloramphenicol acetyl  
transferase (CAT) gene sequence, the mouse  $\beta^{\text{maj}}$ -globin  
poly(A) site, the neomycin gene sequence (Neo), and  
sufficient adenoviral DNA to allow for overlap  
35 recombination.

The reporter vect r was used to generate AdCMV-  
CATNeo, AdCMV-CATN o-dlE3 (AdCMV-CATNeo + pAd70-100dlE3)

to be incorporated by reference and set forth in its entirety here in.

While this invention has been described with emphasis upon preferred embodiments, it will be obvious  
5 to those of ordinary skill in the art that the preferred  
embodiments may be varied. It is intended that the  
invention may be practiced otherwise than as specifically  
described herein. Accordingly, this invention includes  
all modifications encompassed within the spirit and scope  
10 of the appended claims.



## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Wickham, Thomas J.  
Falck-Pedersen, Eric  
Roelvink, Petrus W.  
Bruder, Joseph T.  
Gall, Jason  
Kovesdi, Imre
- (ii) TITLE OF INVENTION: CHIMERIC ADENOVIRAL FIBER PROTEIN AND  
METHODS OF USING SAME
- (iii) NUMBER OF SEQUENCES: 21
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Leydig, Voit & Mayer, Ltd.
  - (B) STREET: Two Prudential Plaza, Suite 4900
  - (C) CITY: Chicago
  - (D) STATE: Illinois
  - (E) COUNTRY: USA
  - (F) ZIP: 60601
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER: US
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (vii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Kilyk, John, Jr.
  - (B) REGISTRATION NUMBER: 30763
  - (C) REFERENCE/DOCKET NUMBER: 71681
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (312) 616-5600
  - (B) TELEFAX: (312) 616-5700

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 31 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown
- (ii) MOLECULE TYPE: DNA (synthetic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCCGCTCGAG TTGCAGATGA AACGCGCCAG A

31

## (2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 35 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unkn wn

32

## (2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 35 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (synthetic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

AGGGCCCCGGG AGGATCCTTA TTCTTGGGCA ATGTA

35

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 55 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (synthetic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TATGGAGGAT CCAATAAAGA ATCGTTTGTG TTATGTTTCA ACGTGTTTAT TTTTC

55

## (2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 57 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (synthetic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

AATTGAAAAA TAAACACGTT GAAACATAAC ACAAACGATT CTTTATTGGA TCCTCCA

57

## (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 32 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (synthetic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AACCCGGTGT ACCCATATGA TGAAAGCAGC TC

32

33

## (2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 35 bas pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (synthetic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

AATGGATCCT CAGTCATCTT CTCTAATATA GGAAA

35

## (2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 79 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (synthetic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATGGATCCAG TACTTTAATT GCGAATGTCT CCGCGTCCAA AACTAGTTCC ACCTCCACCT 60  
CCGAGTTCAT GGATCAAAT

79

## (2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 5 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Val Pro Arg Gly Ser  
1 5

## (2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 54 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..54

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATG	AAG	CGC	GCA	AGA	CCG	TCT	GAA	GAT	ACC	TTC	AAC	CCC	GTG	TAT	CCA	48
Met	Lys	Arg	Ala	Arg	Pro	Ser	Glu	Asp	Thr	Phe	Asn	Pro	Val	Tyr	Pro	
1				5					10					15		

TAT	GAC															54
Tyr	Asp															

## (2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 18 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met	Lys	Arg	Ala	Arg	Pro	Ser	Glu	Asp	Thr	Phe	Asn	Pro	Val	Tyr	Pro
1				5					10					15	

Tyr Asp

35

## (2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 55 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 1..9

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GCC CAA GAA TAAAGAATCG TTTGTGTTAT GTTCAACGT GTTATTTTT CAATTG 55  
Ala Gln Glu  
1

## (2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 3 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Ala Gln Glu  
1

## (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 62 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CATATGGAGG ATCCAATAAA GAATCGTTTG TGTTATGTTT CAACGTGTTT ATTTTCAAT 60  
TG 62

36

## (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 86 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..45

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 49..84

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GGA GGT GGA GGT GGA ACT AGT TTT GGA CGC GGA GAC ATT CGC AAT	45
Gly Gly Gly Gly Gly Thr Ser Phe Gly Arg Gly Asp Ile Arg Asn	
1 5 10 15	
TAA AGT ACT GGA TTC ATG ACT CTA GAC TTA ATT AAG GAT CC	86
Ser Thr Gly Phe Met Thr Leu Asp Leu Ile Lys Asp	
1 5 10	

## (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 15 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Gly Ala Gly Gly Gly Thr Ser Phe Gly Arg Gly Asp Ile Arg Asn	
1 5 10 15	

## (2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 12 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Ser Thr Gly Phe Met Thr Leu Asp Leu Ile Lys Asp	
1 5 10	

37

## (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 98 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 1..51

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GAA CTC GGA GGT GGA GGT GGA ACT AGT TTT GGA CGC GGA GAC ATT CGC 48  
Glu Leu Gly Gly Gly Gly Gly Thr Ser Phe Gly Arg Gly Asp Ile Arg  
1 5 10 15  
AAT TAACTACTG GATTCATGAC TCTAGACTTA ATTAAGGATC CAATAAA 98  
Asn

## (2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 17 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Glu Leu Gly Gly Gly Gly Gly Thr Ser Phe Gly Arg Gly Asp Ile Arg  
1 5 10 15  
Asn

WHAT IS CLAIMED IS:

1. A recombinant adenovirus comprising (a) a chimeric fiber protein, wherein said chimeric fiber protein comprises a nonnative amino acid sequence, which  
5 comprises a protein binding sequence or a protease recognition sequence, in addition to or in place of a native fiber amino acid sequence, which comprises a protein binding sequence, and (b) a therapeutic gene that is capable of being expressed in a cell to which the  
10 recombinant adenovirus attaches or by which the recombinant adenovirus is internalized and, optionally, (c) a nonnative amino acid sequence, which comprises a trimerization domain, in place of a native fiber amino acid sequence, which comprises a trimerization domain.  
15
2. The recombinant adenovirus of claim 1, wherein said nonnative amino acid sequence in said chimeric fiber protein is a protein binding sequence selected from the group consisting of a protein binding sequence from a  
20 serotype of adenovirus different from the serotype of said recombinant adenovirus, a bispecific protein binding sequence, and a multispecific protein binding sequence.
3. The recombinant adenovirus of claim 1, wherein  
25 said nonnative amino acid sequence in said chimeric fiber protein is a protein binding sequence located at the C-terminus of said chimeric fiber protein.
4. The recombinant adenovirus of claim 1, which  
30 further comprises, a chimeric coat protein comprising a nonnative amino acid sequence that is specific for binding to a protein.
5. The recombinant adenovirus of claim 4, wherein  
35 said amino acid sequence in said chimeric coat protein is specific for binding to the same protein as the nonnative amino acid sequence in said chimeric fiber protein.



6. The recombinant adenovirus of claim 4, wherein said chimeric coat protein is selected from the group consisting of a chimeric penton bas and a chimeric hexon.

5

7. A recombinant baculovirus comprising a chimeric adenoviral fiber gene sequence encoding a chimeric fiber protein, which chimeric adenoviral fiber gene sequence comprises a nonnative gene sequence encoding an amino acid sequence, which is a protein binding sequence, in addition to or in place of a native fiber gene sequence that encodes an amino acid sequence, which is a protein binding sequence.

8. A chimeric adenoviral fiber protein comprising a nonnative amino acid sequence, which is a protein binding sequence or a protease recognition sequence, in addition to or in place of a native fiber amino acid sequence, which is a protein binding sequence, and, optionally, a nonnative amino acid sequence, which is a trimerization domain, in place of a native fiber amino acid sequence, which is a trimerization domain.

9. A viral transfer vector comprising a chimeric adenoviral fiber gene sequence encoding a chimeric fiber protein, which chimeric adenoviral fiber gene sequence comprises a nonnative gene sequence encoding an amino acid sequence that is a protein binding sequence or a protease recognition sequence, in addition to a native fiber gene sequence that encodes an amino acid sequence that is a protein binding sequence.

10. The viral transfer vector of claim 9, wherein said viral transfer vector is an adenoviral transfer vector.

11. The viral transfer vector of claim 10, wherein said nonnative gene sequence encoding an amino acid sequence is a protein binding sequence selected from the group consisting of a protein binding sequence from a serotype of adenovirus different from the serotype of said chimeric adenoviral fiber gene sequence, a bispecific protein binding sequence, and a multispecific binding sequence.
12. The viral transfer vector of claim 10, wherein said native fiber gene sequence has been mutated.
13. The viral transfer vector of claim 12, wherein said native fiber gene sequence has been mutated by insertion of the nonnative gene sequence.
14. The viral transfer vector of claim 10, wherein said native fiber gene sequence is a protein binding sequence and has been moved within the fiber gene sequence such that it is conformationally inaccessible for binding to a protein.
15. The viral transfer vector of claim 14, wherein said native fiber gene sequence is a protein binding sequence and has been moved by insertion of said nonnative gene sequence.
16. A viral transfer vector comprising a chimeric adenoviral fiber gene sequence encoding a chimeric fiber protein, which chimeric adenoviral fiber gene sequence comprises a nonnative gene sequence encoding an amino acid sequence, which is a protein binding sequence or a protease recognition sequence, in place of a native fiber gene sequence that encodes an amino acid sequence, which is a protein binding sequence.

17. The viral transfer vector of claim 16, wherein said viral transfer vector is an ad noviral transfer vector.

5        18. The viral transfer vector of claim 17, wherein said nonnative gene sequence encoding an amino acid sequence is a protein binding sequence selected from the group consisting of a protein binding sequence from a serotype of adenovirus different from the serotype of  
10 said chimeric adenoviral fiber gene sequence, a bispecific protein binding sequence, and a multispecific binding sequence.

15        19. A prokaryotic or eukaryotic expression vector comprising a chimeric adenoviral fiber gene sequence encoding a chimeric fiber protein, which chimeric adenoviral fiber gene sequence comprises a nonnative gene sequence encoding an amino acid sequence, which is a protein binding sequence, in addition to or in place of a  
20 native fiber gene sequence that encodes an amino acid sequence, which is a protein binding sequence.

20. A method of gene therapy comprising administering to a patient in need of gene therapy a  
25 therapeutically effective amount of a recombinant adenovirus of claim 1.

21. A method of gene therapy comprising administering to a patient in need of gene therapy a  
30 therapeutically effective amount of a recombinant adenovirus of claim 2.

22. A method of gene therapy comprising administering to a patient in need of gene therapy a  
35 therapeutically effective amount of a recombinant adenovirus of claim 3.

23. A method of gene therapy comprising administering to a patient in need of gene therapy a therapeutically effective amount of a recombinant adenovirus of claim 4.

5

24. A method of gene therapy comprising administering to a patient in need of gene therapy a therapeutically effective amount of a recombinant adenovirus of claim 5.

10

25. A method of gene therapy comprising administering to a patient in need of gene therapy a therapeutically effective amount of a recombinant adenovirus of claim 6.

1/15





FIG. 1

2/15




**Fiber start** **NdeI**  
ATGAAGCGCGCAAGACCGTCTGAAGATACCTTCAACCCCGTGTATCCATATGAC  
►MetLysArgAlaArgProSerGluAspThrPheAsnProValTyrProTyrAsp

(SEQ ID NO:9)

**FIG. 2**  
SUBSTITUTE SHEET (RULE 26)

3/15  
**p193 Ad5 Nd I/SalI**  
(7.29 kB)




**MunI**  
GCCCAAGAATAAAGAATCGTTTGTGTTATGTTTCAACGTGTTTATTTTCAATTG  
▶AlaGlnGlu...

(SEQ ID NO: 11)

**FIG. 3**  
SUBSTITUTE SHEET (RULE 26)

4/15




**FIG. 4**

**SUBSTITUTE SHEET (RULE 26)**



5/15

**p193 FC (F-)**  
(5.604 kB)




**NdeI   BamHI** **MunI**  
CATATGGAGGATCCAATAAAGAATCGTTTGTGTTATGTTTCAACGTGTTTATTTTCAATTG

(SEQ ID NO: 13)

**FIG. 5**  
SUBSTITUTE SHEET (RULE 26)

6/15


**p193 FC (F2)**  
**7299 bp**



**FIG. 6**  
**SUBSTITUTE SHEET (RULE 26)**

7/15


**pGBS.59-100 (F2)**  
**17,269 bp**



**FIG. 7**

**SUBSTITUTE SHEET (RULE 26)**

8/15




**FIG. 8**

SUBSTITUTE SHEET (RULE 26)

9/15


**p193 FC (F3)**  
**6510 bp**



**FIG. 9**  
**SUBSTITUTE SHEET (RULE 26)**

10/15


**pGBS.59-100 (F3)**  
**16,474 bp**



**FIG. 10**

**SUBSTITUTE SHEET (RULE 26)**


11/15



**FIG. 11**

SUBSTITUTE SHEET (RULE 26)

12/15  
**p193 FC (HSF; RGD)**  
 (7.14 kB)



**SpeI** **ScaI**  
 GGAGGTGGAGGTGGAAGTAGTTTTGGACGCGGAGACATTCGCAATTAAAGTACTGGATTCATGACT  
 ►GlyGlyGlyGlyGlyThrSerPheGlyArgGlyAspIleArgAsn...SerThrGlyPheMetThr

**BamHI**  
 CTAGACTTAATTAAGGATCC  
 LeuAspLeuIleLysAsp


(SEQ ID NO: 14)

**FIG. 12**  
 SUBSTITUTE SHEET (RULE 26)



13/15

**pGBS.59-100 (HSF;RGD)**  
(17.106 kB)




(SEQ ID NO: 17)

**FIG. 13**  
SUBSTITUTE SHEET (RULE 26)

14/15

**pAcSG2 (HSF;RGD)**  
(7095 bp)




**SpeI** GGAGGTGGAGGTGGAAGTAGTTTGGACGCGGAGACATTCGCAATTAAAGTACTGGATTCATGACT  
 ▶ GlyGlyGlyGlyGlyThrSerPheGlyArgGlyAspIleArgAsn...SerThrGlyPheMetThr

**BamHI**  
 CTAGACTTAATTAAGGATCC  
 LeuAspLeuIleLysAsp

(SEQ ID NO: 14)

**FIG. 14**  
 SUBSTITUTE SHEET (RULE 26)

15/15



**FIG. 15**  
SUBSTITUTE SHEET (RULE 26)

## INTERNATIONAL SEARCH REPORT

Internat'l Application No

PCT/US 96/01957

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 C12N15/62 C12N15/86 C07K14/075 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,94 10323 (IMP CANCER RES TECH ;SPOONER ROBERT ANTHONY (GB); EPEKETOS AGAMEMN) 11 May 1994	1-3,8-19
Y	see page 26, line 11 - page 27, line 25; claims 1-23	4-7
Y	--- VIROLOGY, NOV 1991, 185 (1) P365-76, UNITED STATES, XP002008215 NOVELLI A ET AL: "Deletion analysis of functional domains in baculovirus-expressed adenovirus type 2 fiber." see the whole document	7
Y	--- WO,A,94 17832 (SCRIPPS RESEARCH INST ;NEMEROW GLEN R (US); WICKHAM THOMAS J (US)) 18 August 1994 see claims 1-20 --- -/-	4-6

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*A\* document member of the same patent family

Date of the actual completion of the international search

11 July 1996

Date of mailing of the international search report

23.07.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+ 31-70) 340-3016

Authorized officer

Gurdjian, D

# INTERNATIONAL SEARCH REPORT

Internat'l Application No

PCT/US 96/01957

C.(Continuation) D CUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VIROLOGY, AUG 1 1994, 202 (2) P782-95, UNITED STATES, XP002008216 KARAYAN L ET AL: "Oligomerization of recombinant penton base of adenovirus type 2 and its assembly with fiber in baculovirus-infected cells." see the whole document ---	1
A	VIROLOGY, vol. 185, no. 2, 1991, pages 758-767, XP002008217 HONG J.S. ET AL.: "The amino terminus of the adenovirus fiber protein encodes the nuclear localization signal" see the whole document ---	3
P,X	J VIROL, MAY 1995, 69 (5) P2850-7, UNITED STATES, XP002008218 STEVENSON SC ET AL: "Human adenovirus serotypes 3 and 5 bind to two different cellular receptors via the fiber head domain." see the whole document ---	1,2,8,9, 19
P,X	WO,A,95 05201 (GENETIC THERAPY INC ;MCCLELLAND ALAN (US); STEVENSON SUSAN C (US)) 23 February 1995 see claims 1-43 -----	1,2,8,9, 19

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/01957

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9410323	11-05-94	EP-A- 0672158 GB-A- 2286593	20-09-95 23-08-95
WO-A-9417832	18-08-94	AU-B- 6133394	29-08-94
WO-A-9505201	23-02-95	EP-A- 0696206	14-02-96